

2017 COA Key West Oral Shell Auction - Bidding List

| Lot | Taxon / Size   | Winning Bid (\$) |
|-----|--|------------------|
| 1   | <i>Nodipecten fragosus</i> (Conrad, 1849) 169.8 purple /113.6yellow/<br>83.8 yellow / 81.7 mm high knobs R7D11.R9D19.R9D20.R9D18 | 500              |
| 2   | <i>Spondylus americanus</i> Hermann, 1781 Purple 191.0/White 189.0mm   | 500              |
| 3   | <i>Scaphella dubia</i> (Broderip, 1827)f. <i>kieneri</i> Clench, 1956<br>160.3/146.8/103.8 mm                                    | 180              |
| 4   | <i>Scaphella junonia butleri</i> Clench, 1953 136.0mm  | 240              |
| 5   | <i>Scaphella junonia</i> (Lamarck, 1840) f <i>johnstoneae</i> Clench, 1953 143.8mm   | 460              |
| 6   | <i>Puntacteon eloiseae</i> (Abbott, 1973) 32.3 / 30.1 /28.3 mm   | 450              |
| 7   | <i>Chama lazarus</i> Linnaeus, 1758 Hypotypes 72.3 / 71.3 / 58.7 / 53.9 mm   | 180              |
| 8   | <i>Bathybembix humboldti</i> Rehder, 1971 53.8mm   | 160              |
| 9   | <i>Tenorioconus aurantius</i> (Hwass in Bruguiere, 1792) 53.0mm  | 400              |
| 10  | <i>Conus bandanus</i> (Hwass in Bruguiere, 1792)<br>[' <i>Conus vidua</i> Reeve, 1843 f. <i>mozoi</i> Melvin, 1980"] 49.2mm      | 130              |
| 11  | <i>Conus barthelemyl</i> (Bemardi 1861) 70.6mm & 57 8mm  | 210              |
| 12  | <i>Tenorioconus cedonulli</i> (Linne, 1767) 49.6mm   | 180              |
| 13  | <i>Tenorioconus cedonulli</i> (Linne, 1767) f. <i>holemani</i> 49.0mm  | 260              |
| 14  | <i>Conus crocatus thailandis</i> da Motta, 1978 69 0mm   | 360              |
| 15  | <i>Conus excelsus</i> G.B. Sowerby III, 1908 76.9mm  | 640              |
| 16  | <i>Conus marmoreus</i> Linnaeus, 1758 59.1mm orange  | 140              |
| 17  | <i>Profundiconus profundorum</i> (Kuroda, 1956) 72.0mm   | 680              |
| 18  | <i>Kurodaconus stupella</i> (Kuroda, 1956) 72.4 / 58 6 mm  | 420              |
| 19  | <i>Conus thomae</i> (Gmelin, 1791) 79.2mm  | 120              |
| 20  | <i>Latiaxis mawae</i> (J.E.Gray in Griffith & Pidgeon, 1834) 41.4 /50.0mm  | 120              |
| 21  | <i>Latiaxis pilsbryi</i> Hirase, 1908 44.7 / 40.0 mm   | 230              |
| 22  | <i>Babelomurex santacruzensis</i> (Emerson & D'Attilio, 1970) 35mm   | 1050             |
| 23  | <i>Hirtomurex teramachii</i> (Kuroda, 1959) 48.9mm   | 180              |
| 24  | <i>Ovatipsa rashleighana</i> (Melvill, 1888) 21.8mm  | 600              |
| 25  | <i>Zoila rosselli</i> Cotton, 1948 61.4mm  | 340              |
| 26  | <i>Luria tessellata</i> (Swainson, 1822) 41.8mm  | 1220             |
| 27  | <i>Luria tessellata</i> (Swainson, 1822) 34.5/33.6 mm  | 700              |

2017 COA Key West Oral Shell Auction - Bidding List

| Lot | Taxon / Size  | Winning Bid (\$) |
|-----|---|------------------|
| 28  | <i>Cirsotrema rugosum</i> (Kuroda & Ito, 1961) 94.8 / 62.0mm  | 260              |
| 29  | <i>Sthenorytis pernobilis</i> (Fischer & Bemardi, 1857) 50.8 / 27.2 mm  | 1400             |
| 30  | <i>Fusinus meyerl</i> (Dunker, 1869) 210.2mm  | 160              |
| 31  | <i>Austroharpa exquisita</i> I redale. 1931 24.0mm  | 400              |
| 32  | <i>Austroharpa loisae</i> Rehder, 1930 23 4mm   | 900              |
| 33  | <i>Austroharpa punctata</i> (Verco, 1896) 38.4mm  | 550              |
| 34  | <i>Harpa gracilis</i> Broderip & G. B. Sowerby I, 1829 22.3mm   | 175              |
| 35  | <i>Harpa major</i> Roding, 1798 f. <i>conoidalis</i> Lamarck, 1822 of authors   | 100              |
| 38  | <i>Morum amabile</i> Shikama, 1973 35.3 / 35.0 / 31.4 mm  | 200              |
| 37  | <i>Morum brunni</i> (Powell, 1958) 37.5mm Hypotype  | 1300             |
| 38  | <i>Morum exquisitum</i> (A. Adams & Reeve, 1858) 34.5mm   | 7000             |
| 39  | <i>Morum grande</i> (A. Adams, 1855) 83.6mm   | 150              |
| 40  | <i>Morum joelgreenei</i> , Emerson, 1981 54.8 / 50.0 / 33.0 mm  | 200              |
| 41  | <i>Morum kurzi</i> Petuch, 1979 24.6mm  | 250              |
| 42  | <i>Morum macandrewi</i> (G.B. Sowerby III, 1889) 34.0 / 27.4 mm   | 600              |
| 43  | <i>Morum matthewsi</i> Emerson, 1967 28 0mm   | 125              |
| 44  | <i>Morum lindae</i> Petuch, 1987 39.9/35.7 mm   | 350              |
| 45  | <i>Morum ponderosum</i> (Hanley, 1858) 36.5 /29.0 mm.<br>The 29.0mm shell was illustrated in Hawaiian Shell News Feb., 1986 | 400              |
| 46  | <i>Morum ninomyai</i> Emerson, 1981 45.9 / 44.0 mm  | 125              |
| 47  | <i>Morum praeclarum</i> Melvill, 1919 36.8 / 30.8 mm  | 175              |
| 48  | <i>Morum uchiyamai</i> Kuroda & Habe 1961 63 7mm  | 175              |
| 49  | <i>Morum teramachii</i> Kuroda & Habe, 1961 54.4 / 54.2 mm  | 80               |
| 50  | <i>Morum veleroae</i> Emerson, 1968 34.5mm  | 10000            |
| 51  | <i>Morum watanabei</i> Kosuge, 1981 42.3 / 37.5 mm  | 85               |
| 52  | <i>Gaza olivacea</i> Quinn, 1992 40.3mm   | 75               |
| 53  | <i>Timbellus bednalli</i> (Brazier, 1878) 78.6/68.9/55.0/40.9mm   | 600              |
| 54  | <i>Favartia</i> ( <i>Murexiella</i> ) <i>bojadorensis</i> (Locard, 1897) 48.4mm   | 350              |
| 55  | <i>Ceratostoma burnetii</i> (A. Adams & Reeve, 1849) 103.3mm & 100.8mm  | 225              |

2017 COA Key West Oral Shell Auction - Bidding List

| Lot | Taxon / Size   | Winning Bid (\$) |
|-----|--|------------------|
| 56  | Homalocantha dovpeledi Houart, 1982 56.8mm                               | 400              |
| 57  | Pterynotus elongatus (Lightfoot, 1786) 100.8 / 89.0 / 47.5 mm            | 75               |
| 58  | Ceratostoma foliatum (Gmelin , 1791) 76.7mm/Albino & 91.6mm              | 550              |
| 59  | Naquetia fosteri D'Attilio & Hertz, 1987 86.4mm                          | 750              |
| 60  | Favartia hidalgoi (Crosse, 1869) 36.2mm                                  | 160              |
| 61  | Chicoreus loebbeckei (Swainson, 1833) 83.4mm                             | 1700             |
| 62  | Purpurellus pinniger (Broderip, 1833) 54 4 /52.9 / 46.4 mm               | 180              |
| 63  | Timbellusus phyllopterus (Lamarck, 1822) 79.6/69.2 mm                    | 1050             |
| 64  | Boreotrophon candelabrum (Reeve, 1848) 48.0mm                            | 150              |
| 65  | Austrotrophon catalinensis (I. Oldroyd, 1927) 88.2mm                     | 3100             |
| 66  | Scabrotrophon emphaticus (Habe & Ito, 1965) 48.6mm                       | 170              |
| 67  | Pagodula fraseri (Knudsen, 1956) 43.8mm                                  | 60               |
| 68  | Stramonistrophon plicatus (Lightfoot, 1786) 61.9mm                       | 660              |
| 69  | Trophonella longstaffi (E.A. Smith, 1907) 40.8mm                         | 3000             |
| 70  | Nipponotrophon stuarti (E.A. Smith, 1880) 57.4mm                         | 380              |
| 71  | Fossil - Ecphora quadricostata (Say, 1824) 100.5mm                       | 360              |
| 72  | Vullietoliva foxi (Stingley, 1984) 29.0mm                                | 175              |
| 73  | Equichlamys bifrons (Lamarck, 1819) 128.2mm                              | 100              |
| 74  | Bractechlamys corallinoides (d'Orbigny, 1840) 56.7mm                     | 225              |
| 75  | Mirapecten cranmerorum (Waller, 1986) 68.6mm                             | 200              |
| 76  | Mirapecten cranmerorum (Waller, 1986) 55.3mm                             | 175              |
| 77  | Semipallium diana (Crandall, 1979) 46.3 / 52.1 /45.3 mm                  | 140              |
| 78  | Anguipecten lamberti (Souverbie in Souverbie & Montrouzier, 1874) 70.6mm | 1200             |
| 79  | Nodipecten magnificus (G.B. Sowerby I, 1835) 194.0 / 124.4 mm            | 1600             |
| 80  | Mirapecten rastellum (Lamarck, 1819) 54.8/45.8 mm Red Sea form           | 75               |
| 81  | Annachlamy reevei (A. Adams & Reeve, 1850) 51.3mm                        | 275              |
| 82  | Annachlamy reevei (A. Adams & Reeve, 1850) 40.2mm                        | 175              |
| 83  | Gloriapallium speciosum (Reeve, 1853) 56.7/55.1 mm                       | 90               |
| 84  | Mimachlamys townsendi (Sowerby III, 1895) 182.9mm                        | 120              |

2017 COA Key West Oral Shell Auction - Bidding List

| Lot | Taxon / Size  | Winning Bid (\$) |
|-----|---|------------------|
| 85  | <i>Entemnotrochus rumphii</i> (Schepman, 1879) 196.5mm  | 1900             |
| 86  | <i>Pleioptygma helenae</i> (Radwin & Bibbey, 1972) 88.0mm   | 280              |
| 87  | <i>Spondylus gloriosus gloriosus</i> Dali, Bartsch & Rehder, 1938<br>Yellow-96mm, Orange- 114.3mm, Purple-125.7mm & White-115.7mm | 1300             |
| 88  | <i>Mikadotrochus salmianus</i> (Rolle, 1899) 105.3mm  | 900              |
| 89  | <i>Tibia insulaechorab</i> Roding, 1798 215.7mm   | 170              |
| 90  | <i>Spondylus senegalensis</i> Schreibers, 1793<br>Two Hypotypes 80.0 / 76.7 / 74.2 / 72.6 / 60.6 mm                               | 1250             |
| 91  | <i>Lambis pilsbryi</i> Abbott, 1961 183.2mm   | 150              |
| 92  | <i>Ophioglossolambis violacea</i> (Swainson, 1821) 127.8mm  | 400              |
| 93  | <i>Angaria vicdani</i> Kosuge, 1980 75.4/74.7 mm  | 250              |
| 94  | <i>Bolma girgyllus</i> (Reeve, 1861) 75.5 / 73.0 mm   | 200              |
| 95  | <i>Altivasum flindersi</i> (Verco, 1914) 156.6mm  | 700              |
| 96  | <i>Cymbiola cymbiola</i> Gmelin, 1791 71.5mm  | 125              |
| 97  | <i>Festilyria duponti</i> Weaver, 1968 107.2mm  | 550              |
| 98  | <i>Amoria grayi kawamurai</i> Habe, 1975 112.2mm  | 2500             |
| 99  | <i>Voluta ebraea</i> Linnaeus, 1758 202.4mm   | 130              |
| 100 | <i>Volutoconus mcmichaeli</i> Habe & Kosuge, 1966 107.5mm   | 1000             |
| 101 | <i>Volutoconus grossi</i> (Iredale, 1927) 130.8mm   | 700              |
| 102 | <i>Volutoconus hargreavesi</i> (Angas, 1872) 114.5mm  | 500              |
| 103 | <i>Volutoconus daisyae</i> Weaver, 1967 93.3mm  | 800              |
| 104 | <i>Volutoconus bednalli</i> (Brazier, 1878) 141.7 / 107.5 mm  | 1000             |
| 105 | <i>Lyria lyraeformis</i> (Swainson, 1821) 132.3/119.8 mm Somalia  | 800              |
| 106 | <i>Lyria lyraeformis</i> (Swainson, 1821) 105.7 / 91.0 mm Kenya   | 650              |
| 107 | <i>Voluta musica</i> Linnaeus, 1758 Seven forms 40mm to 80mm  | 300              |
| 108 | <i>Provocator mirabilis</i> (Finley, 1926) 151.7 / 82.4 mm  | 300              |
| 109 | <i>Cymbiolacca perplicata</i> Hedley, 1902 83.4 / 63.2 mm   | 800              |
| 110 | <i>Voluta polypleura</i> Crosse, 1876 87.3mm  | 300              |
| 111 | <i>Festilyria ponsonbyi</i> (E.A. Smith, 1901) 82.4mm   | 900              |
| 112 | <i>Cymbiola thatcheri</i> (McCoy, 1868) 131.2mm   | 3000             |

2017 COA Key West Oral Shell Auction - Bidding List

| Lot | Taxon / Size   | Winning Bid (\$) |
|-----|--|------------------|
| 113 | <i>Cymbiola thatcheri</i> (McCoy, 1868) 108.0mm  | 450              |
| 114 | 47 species Conidae illustrated in Wall's Cone Shells book  | 750              |
| 115 | 24 species Costellariidae illustrated in Kaicher's catalog and Pechar, Prior & Parkinson's book  | 250              |
| 116 | 60 species Mitridae illustrated in Sally Kaicher's catalog                                       | 250              |
| 117 | <i>Tonna luteostoma</i> (Kuster, 1857) 197.3mm   | 100              |
| 118 | <i>Tonna tankervillei</i> (Henley, 1860) 199.7mm   | 350              |
| 119 | World Record Size shells: Sold as a single lot   | 3500             |
|     | <i>Heliacus stramineus</i> (Gmelin, 1791) 43.9mm WRS Current record is 39 48mm                   |                  |
|     | <i>Vexillum sanguisuga</i> (Linnaeus, 1758) 56.2mm WRS Current record is 42.3mm                  |                  |
|     | <i>Amalda vernelei</i> (G.B. Sowerby II, 1859) 111.4mm WRS Current record is 106.2mm             |                  |
|     | <i>Talochlamys abscondita</i> (P. Fischer in Locard, 1898) 70 3mm, WRS Current record is 43.24mm |                  |
|     | <i>Bractechlamys nodulifera</i> (G.B.Sowerby II, 1842) 57.9mm, WRS Current record is 52.7mm      |                  |
|     | <i>Tudivasum kurzi</i> (MacPherson, 1963) 94.7 mm WRS Current record is 93.9mm                   |                  |
|     | <i>Cymbiola rutila</i> (Broderip, 1826) 137.4mm WRS Current record is 137.3mm                    |                  |

Total (119 lots ) - 81860